

ALLERGENIC EXTRACT

Suggested Dosage Schedule

Veterinarian: _____

Clinic: _____

Phone: _____

Pet's Name/Lab #: _____

Rx Vial #3 (C – Red) Serial #: _____

Special Instructions: _____

**** Clients/Owners administering injections should contact their veterinarian with questions and to re-order vials.**

Maintenance Refill Schedule

The dose and frequency of maintenance injections depends on the progress of controlling the symptoms. The first injection from a new refill vial is 0.1 ml. The maintenance dose is the amount and frequency given that controls symptoms, typically 0.5 ml to 1.0 ml of 1:60w/v*. Do not exceed 1.0 ml dose. Maintenance doses are given monthly with frequency adjustments during peak allergy season.

15 Days After Last Shot

	DAY	DOSE	DATE
VIAL C (Red) (1-60 w/v)	1	0.1ml	
	8	0.5ml	
	15	1.0ml	
	45	1.0ml	
	Monthly	1.0ml	
	Monthly	1.0ml	
	Monthly	1.0ml	
**Re-order Maintenance Vials	Monthly	1.0ml	
	Monthly	1.0ml	
	Monthly	1.0ml	

1. At the discretion of the veterinarian, a responsible owner may administer the injections. This injection schedule is a guideline. Some animals may require modifications to the schedule to suit their individual needs. Please contact your veterinarian.
2. Hyposensitization vials should be refrigerated upon arrival. Refrigeration is not required for shipping. However, vials should be refrigerated when they arrive. **DO NOT FREEZE.**
3. Injections are administered subcutaneously using sterile needles and syringes. **NEVER INJECT INTRAVENOUSLY.** Before injecting, withdraw plunger slightly. If blood appears in syringe, reinsert in another site.
4. Animals should be observed for 30 minutes following injections for possible reactions. Minor reactions to the hyposensitization may occur and are no need for alarm. Signs of such reactions may be increased itching, vomiting, diarrhea, and hives and may occur up to 24 hours past injection. In the event of one or more of these symptoms occurring, temporarily stop further injections. **DO NOT DISCONTINUE ITS USE ENTIRELY.** Call you veterinarian as soon as possible. Your veterinarian may call Pet Preferred Diagnostics for a schedule adjustment. (In most cases a slight schedule adjustment is all that is necessary to continue successful treatment).
5. When possible, the animal should be taken off steroids throughout the entire period of hyposensitization. The most critical period of steroid withdrawal is during vial A. If this is not possible, the lowest possible dose of an oral steroid should be used.
6. Pet Owners: For best results, adhere to the recommended schedule as closely as possible. Immunotherapy may be needed for the lifetime of the animal.
7. When administering a two-vial treatment set, an injection should be given from each vial concurrently at different sites.
8. See package insert for additional information.